



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

**2714 [June, 1918]. Proposed by H. R. HOWARD, University of St. Francis Xavier's College, Nova Scotia.**

A shuffled pack of  $2(p + q)$  cards contains  $2p$  honors. Show that the chance of securing exactly half the honors in taking half the pack is  $[F(p, q)]^2 \div F(2p, 2q)$ , where  $F(p, q)$  denotes the number of different sets of  $p$  cards which can be selected from  $(p + q)$  cards.

Show also that if one honor is removed from the pack, the chance is not thereby affected. Is this true for the chance of getting any other assigned number of honors?

#### SOLUTION BY THE PROPOSER.

We have to find first the number of ways in which we can get exactly  $p$  honors in  $(p + q)$  cards.

We can obviously choose our " $p$ " honors in  $(2p)!/(p!p!)$  ways and the  $q$  other cards in  $(2q)!/(q!q!)$  ways.

Hence, we can effect the required division in  $[(2p)! (2q)! / (p!)^2 (q!)^2]$  ways. Now the number of ways of taking  $(p + q)$  cards from the full pack is  $[2(p + q)]! / (p + q)!^2$ . Thus the chance is

$$\frac{(2p)! (2q)!}{(p!)^2 (q!)^2} \div \frac{[2(p + q)]!}{(p + q)!^2}, \quad (1)$$

i.e.,

$$\frac{(p + q)!^2}{(p!)^2 (q!)^2} \div \frac{(2p + q)!}{(2p)! (2q)!},$$

i.e.,

$$[F(p, q)]^2 \div F(2p, 2q).$$

Now suppose one honor removed from the pack. Then the number of ways of taking  $(p + q)$  cards from the remainder and obtaining exactly  $p$  honors is

$$\frac{(2p - 1)!}{(p - 1)! (p)!} \cdot \frac{(2q)!}{(q!)^2},$$

and the chance is

$$\frac{(2p - 1)!}{(p - 1)! (p)!} \cdot \frac{(2q)!}{(q!)^2} \div \frac{(2p + 2q - 1)!}{(p + q)! (p + q - 1)!}. \quad (2)$$

Dividing (1) by (2) we obtain unity for the quotient and this proves their equality. Let  $x$  be the assigned number of honors. We shall show that the condition that the chances be equal can only be satisfied by  $x = p$ . With the full pack the chance will now be

$$\frac{(2p)!}{x! (2p - x)!} \cdot \frac{(2q)!}{(p + q - x)! (q - p + x)!} \div \frac{(2p + 2q)!}{(p + q)!^2}. \quad (3)$$

With one honor removed the chance will be

$$\frac{(2p - 1)!}{x! (2p - x - 1)!} \cdot \frac{(2q)!}{(p + q - x)! (q - p + x)!} \div \frac{(2p + 2q - 1)!}{(p + q)! (p + q - 1)!}. \quad (4)$$

Dividing (3) by (4) we see that these can only be equal if

$$\frac{p}{2p - x} = 1$$

or  $x = p$ , which proves our statement.

Also solved by H. L. OLSON, A. PELLETIER, and E. E. WITMER.

**2715 [June, 1918]. Proposed by H. R. KINGSTON, University of Manitoba.**

$A', B', C'$  are points on the sides  $BC, CA, AB$ , respectively, of the triangle  $ABC$ , and  $AA', BB', CC'$  are concurrent in  $O$ .  $X, Y, Z$  are the three collinear points in which, by Desargues' theorem, the corresponding sides of the triangles  $ABC$  and  $A'B'C'$  intersect. If  $A'', B'', C''$  are the vertices of the triangle formed by the lines  $AX, BY, CZ$ , show that  $AA'', BB'', CC''$  are concurrent.